

Numbers and periods

Notes for a talk at the H-C.R.I., 8 February 2006

Chandan Singh Dalawat

We review the construction, starting with the monoid $\mathbf{N}$ of natural numbers, of the ring $\mathbf{Z}$ of integers, of the field $\mathbf{Q}$ of rational numbers, of the locally compact fields $\mathbf{R}$ and $\mathbf{Q}_p$ of real and p -adic numbers (p being a prime number), and of the complete closed fields $\mathbf{C}$ and $\mathbf{C}_p$. We introduce Fontaine's rings of periods and, finally, indicate the need for going beyond them for the purposes of outer galoisian actions.

1. (*Numbers as sums*). Mathematics starts with

DEFINITION 1. — *The monoid $\mathbf{N}$ is the free monoid on one generator, denoted 1.*

We grant that such a monoid exists. The elements of $\mathbf{N}$ are $0, 1, 1 + 1, 1 + 1 + 1, \dots$; they are to be added for example as

$$(1 + 1) + (1 + 1 + 1) = 1 + 1 + 1 + 1 + 1$$

with the requirement that $n + 0 = n$ and $0 + n = n$ for every $n \in \mathbf{N}$. Given any monoid M and an element $m \in M$, there is a unique morphism of monoids $f : \mathbf{N} \rightarrow M$ such that $f(1) = m$.

Summary : $(\mathbf{N}, 1)$ is the initial object in the category of pointed monoids.

2. (*Numbers as products*). Let $\text{End}(\mathbf{N})$ be the additive monoid of endomorphisms of $\mathbf{N}$:

$$(f + g)(n) = f(n) + g(n).$$

The map $e : \text{End}(\mathbf{N}) \rightarrow \mathbf{N}$, $f \mapsto f(1)$ is an isomorphism of monoids; the reciprocal is the morphism $1 \mapsto (1 \mapsto 1)$, i.e. the unique morphism which sends 1 to $\text{Id}_{\mathbf{N}}$.

An endomorphism f is injective if and only if $f(1) \neq 0$, bijective if and only if $f(1) = 1$. As the composite of two injective maps is injective, e can be used to make the set $\mathbf{N}^*$ (the complement of 0 in $\mathbf{N}$) into a monoid, by transport of structure; it is *commutative*.

For $m, n \in \mathbf{N}^*$, we say that m divides n , and write $m|n$, if the endomorphism $1 \mapsto n$ factors via the endomorphism $1 \mapsto m$.

DEFINITION 2. — A number $p \in \mathbf{N}^*$ is called prime if $p \neq 1$ and if the only divisors of p are 1 and p .

The set of prime numbers is denoted by P . The first few primes are 2, 3, 5, 7, 11, ... Multiples of 2 are called even numbers, the others are called odd. In particular, every prime $\neq 2$ is odd.

THEOREM 1. — The map $\mathbf{N}^{(P)} \rightarrow \mathbf{N}^*$ which send f to $\prod_{p \in P} p^{f(p)}$ is an isomorphism, i.e. $\mathbf{N}^*$ is the free commutative monoid on the set P .

In other words, given any commutative monoid M and any map $f : P \rightarrow M$, there is a unique morphism $\mathbf{N}^* \rightarrow M$ which extends f . Moreover, P is the only subset of $\mathbf{N}^*$ with this property.

THEOREM 2. — The monoid $\mathbf{N}^*$ is not finitely generated : P is infinite.

Proof (Euclid) : Suppose that P were finite, and consider the number $n = 1 + \prod_{p \in P} p$, which is > 1 . If n is prime, we get a contradiction because P does not contain n . If n is not a prime, we also get a contradiction, because it does not have any prime divisor : no $p \in P$ is a divisor of n .

Summary : $\mathbf{N}^*$ is the monoid of injective endomorphisms of the monoid $\mathbf{N}$; it is free commutative; the elements of its basis are called prime numbers.

3. (Numbers as differences). If $f : \mathbf{N} \rightarrow M$ is a morphism of monoids such that $f(1)$ has an (additive) inverse, then $f(n)$ has an inverse for every $n \in \mathbf{N}$. Conversely, if $f(n)$ is invertible for some $n \neq 0$, then $f(1)$ is invertible.

The trouble is that there is no $n \in \mathbf{N}$ such that $n + 1 = 0$, i.e. the difference $0 - 1$ does not exist. Nor does the difference $8 - 9$, which is essentially the same. The way to make them exist is to declare these differences $m - n$ ($m, n \in \mathbf{N}$) themselves as being new numbers and to identify any two differences which are essentially the same.

Formally, consider the equivalence relation $\sim$ on the monoid $\mathbf{N} \times \mathbf{N}$ of all pairs, defined as

$$(m, n) \sim (m', n') \Leftrightarrow m + n' = m' + n.$$

Then the class of $(0, 1)$ is the same as the class of $(8, 9)$, and so on; it is denoted -1 . The pairs $(1, 0)$, $(9, 8)$, etc., have the same class, which is identified with 1. Thus the equivalence classes consist of

$$\dots, -3, -2, -1, 0, 1, 2, 3, \dots$$

and we have obtained the group $\mathbf{Z}$ of integers as the group of differences of the monoid $\mathbf{N}$

THEOREM 3. — *The group $\mathbf{Z}$ is free on one generator; 1 and -1 are the only generators.*

Summary : $\mathbf{N} \rightarrow \mathbf{Z}$ is the group of differences of $\mathbf{N}$, i.e. the initial object in the category of maps of the monoid $\mathbf{N}$ into groups.

4. (*Numbers as products, bis*). Consider the ring $\text{End}(\mathbf{Z})$ of endomorphisms of the group $\mathbf{Z}$. The map $\text{End}(\mathbf{Z}) \rightarrow \mathbf{Z}, f \mapsto f(1)$ is an isomorphism of additive groups, allowing us to make the commutative group $\mathbf{Z}$ into a ring; extending the multiplication from $\mathbf{N}$ to $\mathbf{Z}$. This ring is commutative and integral, i.e., if $xy = 0$ ($x, y \in \mathbf{Z}$), then $x = 0$ or $y = 0$. Moreover, the sum and the product of two positive numbers is again positive.

THEOREM 4. — *The ring $\mathbf{Z}$ is principal, i.e., every ideal can be generated by one element.*

Given any ring A , there is a unique homomorphism of rings $\mathbf{Z} \rightarrow A$, a fact which is expressed by saying that the ring $\mathbf{Z}$ is the initial object in the category of rings.

THEOREM 5. — *For every prime p , the ideal $p\mathbf{Z}$ is prime, indeed maximal. Conversely, every maximal ideal of $\mathbf{Z}$ is generated by a prime number.*

Notice that 0 is the only prime ideal of $\mathbf{Z}$ which is not maximal. For every prime p , the field $\mathbf{Z}/p\mathbf{Z}$ is denoted $\mathbf{F}_p$.

Summary : The ring $\mathbf{Z}$ is the ring of endomorphisms of the (commutative) group $\mathbf{Z}$. It is principal and the maximal ideal correspond to prime numbers.

5. (*Numbers as fractions*). We notice that fractions do not exist in $\mathbf{Z}$: there is no x such that $2x = 1$. In order to remedy this, we proceed as with difference : on the set $\mathbf{Z} \times \mathbf{Z}^*$ of pairs (m, n) ($n \neq 0$), we define an equivalence relation $\sim$ as

$$(m, n) \sim (m', n') \Leftrightarrow mn' = m'n.$$

The equivalence class of (m, n) is written m/n ; they are to be added and multiplied as

$$\frac{m}{n} + \frac{m'}{n'} = \frac{mn' + m'n}{nn'}, \quad \frac{m}{n} \cdot \frac{m'}{n'} = \frac{mm'}{nn'}.$$

These classes form a field $\mathbf{Q}$, called the field of fractions of the (integral) ring $\mathbf{Z}$.

Consider the multiplicative group $\mathbf{Q}^\times$ of the field $\mathbf{Q}$, and its subgroup $\mathbf{Z}^\times$, which happens to be the torsion subgroup. One identifies $\mathbf{P}$ with its image in $\mathbf{Q}^\times/\mathbf{Z}^\times$.

THEOREM 6. — *The map $\mathbf{Z}^{(\mathbf{P})} \rightarrow \mathbf{Q}^\times/\mathbf{Z}^\times$ which sends f to $\prod_{p \in \mathbf{P}} p^{f(p)}$ is an isomorphism, i.e. $\mathbf{Q}^\times/\mathbf{Z}^\times$ is the free $\mathbf{Z}$ -module on the set $\mathbf{P}$.*

In other words, $\mathbf{Q}^\times/\mathbf{Z}^\times$ is the group of “differences” of the monoid $\mathbf{N}^*$.

COROLLARY. — *The map $\{1, -1\} \times \mathbf{Z}^{(\mathbf{P})} \rightarrow \mathbf{Q}^\times$ which sends (ε, f) to $\varepsilon \prod_{p \in \mathbf{P}} p^{f(p)}$ is an isomorphism of $\mathbf{Z}$ -modules.*

Sometimes it is more convenient to use the isomorphism which sends -1 to -1 and e_p to p^* for every prime p , where $(e_p)_{p \in \mathbf{P}}$ is the canonical basis of $\mathbf{Z}^{(\mathbf{P})}$ and

$$2^* = 2, \quad p^* = p^{\frac{p-1}{2}} \quad (p \neq 2).$$

Summary : $\mathbf{Z} \rightarrow \mathbf{Q}$ is the field of fractions of $\mathbf{Z}$, i.e. the initial object in the category of injective homomorphisms of the ring $\mathbf{Z}$ into fields. $\mathbf{N}^* \rightarrow \mathbf{Q}^\times/\mathbf{Z}^\times$ is the “group of differences” of $\mathbf{N}^*$.

6. (*Numbers as limits*). We notice that there are sequences $(x_n)_n$ of rational numbers which do not have a limit in $\mathbf{Q}$, although $|x_m - x_n|$ can be made as small as we please by taking m, n large enough. For example,

$$x_n = \frac{1}{0!} + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \cdots + \frac{1}{n!}$$

In order to make such limits exist, we consider the space of all fundamental sequences, i.e. sequences $(x_n)_n$ of rational numbers such that $|x_n - x_m|$ can be made as small as we please by taking m, n large enough. They can be added and multiplied, so they form a ring. Those fundamental sequences which tend to 0, i.e. such that $|x_n|$ is as small as we please for n large enough, form a maximal ideal of this ring. The quotient field $\mathbf{R}$ has a natural order, hence a uniformity and a topology.

THEOREM 7. — *The space $\mathbf{R}$ is connected, locally connected, locally compact and complete.*

7. (*Numbers as roots*). The polynomial $X^2 + 1$ is irreducible over $\mathbf{R}$ because $x^2 + 1 > 0$ for every $x \in \mathbf{R}$. Adjoining a root of this polynomial, we obtain the field $\mathbf{C} = \mathbf{R}[X]/(X^2 + 1)$ of complex numbers. Something miraculous happens :

THEOREM 8. — *The field $\mathbf{C}$ of complex numbers is algebraically closed.*

The proof rests on the fact that every real number > 0 has a square root, and that every real polynomial of odd degree has a root in $\mathbf{R}$.

The group of $\mathbf{R}$ -automorphisms of $\mathbf{C}$ is generated by the involution $i \mapsto -i$, where i is the image of X in $\mathbf{C}$.

The field $\mathbf{C}$ comes with an absolute value $|\cdot|_\infty$ extending the one on $\mathbf{R}$, which can be used to give it a uniformity, and hence a topology. Another miracle happens :

THEOREM 9. — *The space $\mathbf{C}$ is simply connected, locally compact and complete.*

Let us mention the field $\mathbf{H}$ of quaternions, which is not commutative. It was discovered by Hamilton, who was trying to find an even bigger number system than $\mathbf{C}$. It can now be defined as the unique (4-dimensional) $\mathbf{R}$ -algebra A , other than $\mathbf{M}_2(\mathbf{R})$, such that $A \otimes_{\mathbf{R}} \mathbf{C}$ is $\mathbf{C}$ -isomorphic to $\mathbf{M}_2(\mathbf{C})$. The absolute value $|\cdot|_\infty$ extends to $\mathbf{H}$, with respect to which it is complete.

THEOREM 10. — *The only fields complete with respect to an archimedean absolute value are $\mathbf{R}$, $\mathbf{C}$ and $\mathbf{H}$.*

In fact, these are the only locally compact connected fields. Notice that these three fields are different, as $\mathbf{R}$ is commutative but not algebraically closed, $\mathbf{C}$ is commutative and algebraically closed, whereas $\mathbf{H}$ is not commutative.

Summary : $\mathbf{Q} \rightarrow \mathbf{R}$ is the completion of $\mathbf{Q}$ in the ∞ -adic uniformity. It is connected, locally connected, locally compact and complete. $\mathbf{R} \rightarrow \mathbf{C}$ is a quadratic extension and an algebraic closure, with a chosen 4th root of 1.

8. (*Numbers as limits, bis*). It was noticed at the end of the XIXth century that $|\cdot|_\infty$ is not the only absolute value on $\mathbf{Q}$. For every prime p , there is an absolute value $|\cdot|_p$ for which the sequence

$$x_n = 1 + p + p^2 + p^3 + \cdots + p^n$$

is a fundamental sequence! It is the unique homomorphism $\mathbf{Q}^\times \rightarrow \mathbf{R}^{\times\circ}$ (into the multiplicative group of real numbers > 0) such that $|p|_p = 1/p$ and $|l|_p = 1$ for every prime $l \neq p$ (cf. Corollary to Theorem 6).

THEOREM 11. — *Leaving aside the trivial one, every absolute value on $\mathbf{Q}$ is equivalent to $|\cdot|_\infty$ or to $|\cdot|_p$ for some prime $p \in \mathbf{P}$.*

That none of these absolute values can be neglected is illustrated by the following result (the “product formula”) — almost a tautology — in which $\bar{\mathbf{P}}$ denoted the set $\mathbf{P}$ together with an additional element ∞ :

THEOREM 12. — *For every $x \in \mathbf{Q}^\times$, one has $|x|_v = 1$ for almost all $v \in \bar{\mathbf{P}}$ and $\prod_{v \in \bar{\mathbf{P}}} |x|_v = 1$.*

Indeed, it is sufficient to verify this for $x = -1$ and $x = p$ ($p \in \mathbf{P}$), cf. Corollary to Theorem 6.

There are fundamental sequences with respect to $|\cdot|_p$ which do not converge in $\mathbf{Q}$, for example the sequence $x_n = 2^{5^n}$ for $p = 5$, whose limit would be a primitive 4th root of 1.

As in the construction of real numbers, we can consider the ring of all fundamental sequences; those which converge to 0 form a maximal ideal. The quotient field $\mathbf{Q}_p$ comes with an absolute value $|\cdot|_p$ (extending the one on $\mathbf{Q}$), hence a metric.

THEOREM 13. — *The space $\mathbf{Q}_p$ is locally compact, totally disconnected, and complete.*

The disc $|x|_p \leq 1$ is a compact subring $\mathbf{Z}_p$, with group of units $\mathbf{Z}_p^\times$ the circle $|x|_p = 1$. Another simplification occurs in the multiplicative group (cf. Theorem 6).

THEOREM 14. — *The valuation $v_p : \mathbf{Q}_p^\times \rightarrow \mathbf{Z}$ induces an isomorphism $\mathbf{Q}_p^\times / \mathbf{Z}_p^\times \rightarrow \mathbf{Z}$.*

Choosing a section of v_p , i.e. an element $e_p \in \mathbf{Q}_p^\times$ such that $v_p(e_p) = 1$ (for example $e_p = p$), we get an isomorphism $\mathbf{Z}_p^\times \times \mathbf{Z} \rightarrow \mathbf{Q}_p^\times$. For p odd, the torsion subgroup W of $\mathbf{Z}_p^\times$ is cyclic of order $p-1$; the quotient $\mathbf{Z}_p^\times / W$ is a commutative pro- p -group, i.e. a $\mathbf{Z}_p$ -module; it is free of rank 1; the image of $1 + e_p$ is a basis. What happens for $p = 2$?

What is the archimedean analogue of this theorem? The analogue of the units would be the kernel of the absolute value, i.e. the 0-sphere $\{1, -1\}$ in $\mathbf{R}$, the 1-sphere $\mathbf{S}_1$ in $\mathbf{C}$ and the 3-sphere $\mathbf{S}_3$ in $\mathbf{H}$.

THEOREM 15. — *The maps $(\varepsilon, x) \mapsto \varepsilon e^x$, $\{1, -1\} \times \mathbf{R} \rightarrow \mathbf{R}^\times$, $\mathbf{S}_1 \times \mathbf{R} \rightarrow \mathbf{C}^\times$ and $\mathbf{S}_3 \times \mathbf{R} \rightarrow \mathbf{H}^\times$ are isomorphisms of real-analytic groups.*

In other words, the topological groups $\mathbf{R}^\times / \{1, -1\}$, $\mathbf{C}^\times / \mathbf{S}_1$ and $\mathbf{H}^\times / \mathbf{S}_3$ are 1-dimensional vector spaces over $\mathbf{R}$, with canonical basis $\bar{e}$; they are

the same as the multiplicative group $\mathbf{R}^{\times \circ}$ of reals > 0 .

Further, there is a unique $\pi > 0$ in $\mathbf{R}$ such that $2i\pi\mathbf{Z}$ is the kernel of $e^{(\cdot)} : 2i\pi\mathbf{R} \rightarrow \mathbf{S}_1$, i.e. one has the exact sequence

$$\{0\} \rightarrow 2i\pi\mathbf{Z} \rightarrow 2i\pi\mathbf{R} \rightarrow \mathbf{S}_1 \rightarrow \{1\}.$$

What is the analogue for $\mathbf{S}_3$? What is the liegebra $\mathfrak{s}_3$? What are the groups $\mathbf{C}^\times/\mathbf{R}^\times$, $\mathbf{H}^\times/\mathbf{R}^\times$, $\mathbf{H}^\times/\{1, -1\}$, $\mathbf{S}_1/\{1, -1\}$, $\mathbf{S}_3/\{1, -1\}$?

In all fairness, we should give a similar description of the multiplicative group of all fields complete with respect to the p -adic absolute value, and not just $\mathbf{Q}_p$. Let us leave that aside.

Summary : $\mathbf{Q} \rightarrow \mathbf{Q}_p$ is the completion of $\mathbf{Q}$ in the p -adic uniformity; it is locally compact and totally disconnected. The closed unit disc $\mathbf{Z}_p$ is a compact subring; the quotient $\mathbf{Q}_p^\times/\mathbf{Z}_p^\times$ is $\mathbf{Z}$.

9. (*Numbers as roots and limits, bis*). We saw that there is a drastic reduction in the number of irreducible polynomials when we completed $\mathbf{Q}$ with respect to $|\cdot|_\infty$. The reduction is not so drastic when we complete with respect to $|\cdot|_p$: there are irreducible polynomials of every degree, but essentially only finitely many of them :

THEOREM 16. — *The field $\mathbf{Q}_p$ has only finitely many extensions in each degree.*

For example, the only quadratic extensions of $\mathbf{Q}_5$ are $\mathbf{Q}_5(\sqrt{5})$, $\mathbf{Q}_5(\sqrt{2})$ and $\mathbf{Q}_5(\sqrt{10})$.

Central simple $\mathbf{Q}_p$ -algebras are classified by the group $\mathbf{Q}/\mathbf{Z}$, whereas central simple $\mathbf{R}$ -algebras are classified by the group $\frac{1}{2}\mathbf{Z}/\mathbf{Z}$. There is a very precise local-to-global principle in terms of these groups, of which a special case reads :

THEOREM 17. — *An n^2 -dimensional central simple $\mathbf{Q}$ -algebra A is isomorphic to $\mathbf{M}_2(\mathbf{Q})$ if $A \otimes_{\mathbf{Q}} \mathbf{Q}_p$ is isomorphic to $\mathbf{M}_2(\mathbf{Q}_p)$ for every $p \in \mathbf{P}$; $A \otimes_{\mathbf{Q}} \mathbf{R}$ is then necessarily isomorphic to $\mathbf{M}_2(\mathbf{R})$.*

Let $\overline{\mathbf{Q}}_p$ be an algebraic closure of $\mathbf{Q}_p$; the group $\text{Gal}(\overline{\mathbf{Q}}_p|\mathbf{Q}_p)$, being pro-solvable, is simpler than $\text{Gal}(\overline{\mathbf{Q}}|\mathbf{Q})$, but it is considerably more complicated than the order-2 group $\text{Gal}(\mathbf{C}|\mathbf{R})$. The absolute value $|\cdot|_p$ extends uniquely to $\overline{\mathbf{Q}}_p$. Here is a minor embarassment :

THEOREM 18. — *The field $\overline{\mathbf{Q}}_p$ is not complete (and not locally compact).*

But we can complete $\overline{\mathbf{Q}}_p$ with respect to $|\cdot|_p$ to obtain a field $\mathbf{C}_p$, to which $|\cdot|_p$ extends uniquely, with the same group of values.

THEOREM 19. — *The field $\mathbf{C}_p$ is algebraically closed.*

So we don't need to start all over again, taking an algebraic closure, etc.

Summary : $\overline{\mathbf{Q}}_p$ is an algebraic closure of $\mathbf{Q}_p$. $\mathbf{C}_p$ is the completion of $\overline{\mathbf{Q}}_p$ in the p -adic uniformity; it is algebraically closed.

10. (*Numbers as periods*). Varieties have periods. Let us return briefly to the archimedean world of $\mathbf{R}$ and $\mathbf{C}$ to illustrate this.

A considerable effort has been spent, following Grothendieck's insights, in the search for a universal cohomology theory for varieties over $\mathbf{Q}$. It was noticed that the various known cohomology theories share a number of properties, which led Grothendieck to suspect that there must be a universal cohomology theory — a theory of motives —, of which the known cohomology theories are but various different “avatars”. In other words, the motive of a variety should carry every bit of cohomological information about the variety.

To be specific, let us take a smooth, projective, absolutely connected curve X of genus g over $\mathbf{R}$. Then $H^1(X(\mathbf{C}), \mathbf{Z})$ is a free $\mathbf{Z}$ -module of rank $2g$, so all it knows about X is its genus. However, the $\mathbf{C}$ -space $H^1(X(\mathbf{C}), \mathbf{Z}) \otimes \mathbf{C}$ has a natural filtration which carries much more information about X , and indeed enough information to recover the jacobian J of X . What I want to emphasize is that you need to tensor with $\mathbf{C}$ to get this additional structure.

The miracle is that tensoring with $\mathbf{C}$ suffices for varieties over $\mathbf{R}$. Another way of saying this is that the field $\mathbf{C}$ contains the periods of all smooth projective $\mathbf{R}$ -varieties. This is a miracle because $\mathbf{C}$ was not designed to serve this purpose; it was merely designed to have a square root of -1 .

11. (*Numbers as periods, bis*). Varieties over $\mathbf{Q}_p$ have their periods too. The major difference with the archimedean world is that you have to go way beyond $\mathbf{C}_p$ to get all the periods. Fontaine has pursued relentlessly this quest for the p -adic analogue of $2i\pi$; his solution is to construct several rings carrying many structures which play the role of rings of periods for different kinds of varieties : those with smooth reduction, those with semistable reduction, all (smooth proper) varieties.

The rings B constructed by Fontaine are $\mathbf{Q}_p$ -algebras which carry — among other structures — a representation of $G_p = \text{Gal}(\overline{\mathbf{Q}_p}|\mathbf{Q}_p)$; the invariants B^{G_p} are a field. A p -adic representation E of G_p , i.e. a finite-dimensions vector $\mathbf{Q}_p$ -space with a continuous linear action of G_p , is called B -admissible if the dimension of $(B \otimes_{\mathbf{Q}_p} E)^{G_p}$ (as a vector space over B^{G_p}) is the same as the dimension of E (as a vector space over $\mathbf{Q}_p$). Notable examples of p -adic representations are provided by the p -adic étale cohomology of varieties over $\mathbf{Q}_p$ — the analogue of the singular cohomology of varieties over $\mathbf{R}$ and $\mathbf{C}$.

The rings of Fontaine have proved their utility beyond any doubt. A theorem of Colmez gives an idea of their coherence as p varies : he proves a product formula (cf. Theorem 12) for the periods of abelian varieties having complex multiplications, periods which may be transcendental!

I have chosen to illustrate their utility by taking for B the crystalline ring of Fontaine; B -admissible representations of G_p are called crystalline.

THEOREM (Faltings). — *For a smooth projective variety X having smooth reduction over $\mathbf{Q}_p$, the representation $H^n(X_{\overline{\mathbf{Q}_p}}, \mathbf{Q}_p)$ is crystalline.*

The converse need not hold. Nevertheless, in the light of the Serre-Tate criterion (see below), one has a right to expect it to hold for abelian varieties.

THEOREM (Coleman-Iovita). — *Let A be an abelian $\mathbf{Q}_p$ -variety. If the p -adic representation $H^1(A_{\overline{\mathbf{Q}_p}}, \mathbf{Q}_p)$ is crystalline, then A has abelian reduction.*

If we compare these two theorems with what goes on for l -adic étale cohomology, where l is a prime $\neq p$, the similarity is striking :

THEOREM (Grothendieck). — *For a smooth projective variety X having smooth reduction over $\mathbf{Q}_p$, the representation $H^n(X_{\overline{\mathbf{Q}_p}}, \mathbf{Q}_l)$ is unramified.*

THEOREM (Serre-Tate). — *Let A be an abelian $\mathbf{Q}_p$ -variety. If the l -adic representation $H^1(A_{\overline{\mathbf{Q}_p}}, \mathbf{Q}_l)$ is unramified, then A has abelian reduction.*

Summary : For abelian varieties over a finite extension of $\mathbf{Q}_p$, there is an l -adic and a p -adic criterion for abelian reduction.

12. (Anabelian periods). These criteria of Serre-Tate and Coleman-Iovita fail for curves. Indeed, it is not very difficult to construct smooth

proper absolutely connected curves C over $\mathbf{Q}_p$ which do not have smooth reduction but whose jacobian has abelian reduction. For such a curve C , the l -adic étale cohomology $H^1(C_{\overline{\mathbf{Q}}_p}, \mathbf{Q}_l)$ is unramified and the p -adic étale cohomology $H^1(C_{\overline{\mathbf{Q}}_p}, \mathbf{Q}_p)$ is crystalline, because these groups are the same for C and its jacobian.

Is there a criterion for the smooth reduction of curves? A very striking theorem was proved by Takayuki Oda, who gives an anabelian l -adic criterion. He replaces $H^1(C_{\overline{\mathbf{Q}}_p}, \mathbf{Q}_l)$ by the maximal pro- l -quotient $\pi_1^{(l)}(C_{\overline{\mathbf{Q}}_p})$ of the fundamental group $\pi_1(C_{\overline{\mathbf{Q}}_p})$ of $C_{\overline{\mathbf{Q}}_p}$ (there is a choice of a base point involved; it will turn out to be immaterial). The fundamental group carries a natural outer action of the Galois group, i.e. there is a natural homomorphism $G_p \rightarrow \text{Out } \pi_1^{(l)}(C_{\overline{\mathbf{Q}}_p})$.

THEOREM (Takayuki Oda). — *Let C be a smooth proper absolutely connected $\mathbf{Q}_p$ -curve of genus > 1 . If the outer action $G_p \rightarrow \text{Out } \pi_1^{(l)}(C_{\overline{\mathbf{Q}}_p})$ on the pro- l π_1 is unramified, then C has smooth reduction.*

In fact, his theorem works over any finite extension of $\mathbf{Q}_p$.

Summary : For curves, there is an anabelian l -adic criterion for smooth reduction.

13. (*Anabelian periods, bis*). Let me end with a problem which I would very much like to solve. Anyone who has read this account cannot fail to ask himself : Is there an anabelian p -adic criterion for smooth reduction of curves over $\mathbf{Q}_p$? I happened to mention this to Kazuya Kato, and he said that this problem “goes straight to the heart”.

PROBLEM. — *Give a necessary and sufficient condition on the outer action $G_p \rightarrow \text{Out } \pi_1^{(p)}(C_{\overline{\mathbf{Q}}_p})$ for the (smooth, projective, absolutely connected) $\mathbf{Q}_p$ -curve C (of genus > 1) to have smooth reduction.*

In other words, one should complete the square :

	$l \neq p$	$l = p$
	Serre-Tate	Coleman-Iovita
abelian varieties	(1968, “unramified”)	(1999, “crystalline”)
	Takayuki Oda	???
curves of genus > 1	(1995, “unramified”)	(???)

The problem should be studied not just over $\mathbf{Q}_p$, but over every finite extension thereof.

Summary : Find an anabelian p -adic criterion for smooth reduction of curves.

14 (*Semistable reduction*). Notice that there is already a p -adic criterion for semistable reduction of curves and abelian varieties in terms of Fontaine's semistable ring. In effect, Fontaine has proved that if an abelian K -variety has semiabelian reduction, K being a finite extension of $\mathbf{Q}_p$, then the representation $H^1(A_{\overline{K}}, \mathbf{Q}_p)$ of $\text{Gal}(\overline{K}|K)$ is semistable, $\overline{K}$ being an algebraic closure of K . Later, Breuil proved the converse, at least for $p \neq 2$. Earlier, Deligne and Mumford had proved that for a curve to have semistable reduction, it is necessary and sufficient for its jacobian to have semiabelian reduction.